






The Full Spectrum Protocol – Trial 3 Results:




The third trial was conducted with 16 participants across South Africa, ranging in age from 3 to 55 years old, over a 2-full month period (June–August 2025).

Some participants were able to transition off their pharmaceutical regimen, while others used the protocol in combination with existing treatments. We have noted that outcomes were generally less pronounced in the latter case.

In many instances, the protocol produced remarkable improvements within only two month, offering tangible relief to those families. Though in a few instances, changes were less noticable, highlighting areas for future consideration when in combination with pharmaceuticals.

			AVERAGE
1	On a scale of 0–100%, how would you rate the change in your child's communication skills after the 2 month trial?		48%
2	On a scale of 0–100%, how much improvement have you observed in your child's ability to express their needs and feelings after the 2 month trial?		52%
3	On a scale of 0–100%, how much improvement have you observed in your child's social interaction with family and friends after the 2-month trial?		50%
4	On a scale of 0–100%, how much improvement have you observed in your child's social interactions within school or peer environments after the 2- month trial, compared to before?		50%
5	On a scale of 0–100%, how would you rate the change in your child's emotional regulation after the 2-month trial?		56%

6	On a scale of 0–100%, how would you rate the change in your child's daily living skills and independence after the 2-month trial?	6	55%
7	On a scale of 0–100%, how much improvement have you observed in your child's independence in performing daily tasks after the 2-month trial?	7	51%
8	On a scale of 0–100%, how would you rate the change in your child's sensory and physical behaviours after the 2-month trial?	8	49%
9	On a scale of 0–100%, how much improvement have you observed in your child's participation and engagement with parents and family after the 2-month trial?	9	54%
10	On a scale of 0–100%, how much improvement have you observed in your child's interaction with daily family life after the 2-month trial?	10	62%
11	On a scale of 0–100%, how much improvement have you observed in your child's ability to respond to changes in routine or environment after the 2-month trial?	11	53%

			Slightly less Frequently	Noticeably less frequently	Rarely occurs now	Has Stopped Completely	No Change
12	How frequently does your child experience meltdowns or emotional outbursts after the 2-month trial, compared to before?	 12 20% 35% 29% 8% 8%	20%	35%	29%	8%	8%
			Yes	No	Unsure		
13	Based on your experience, how likely are you to continue using this protocol for your child?	 13 70% 10% 20%	70%	10%	20%		
			Yes	No	Unsure		
14	Based on your experience, how likely are you to recommend this protocol to friends and family?	 14 83% 0% 17%	83%	0%	17%		